Primary duodenal adenocarcinoma of the fourth portion diagnosed using double-balloon enteroscopy and surgically resected: A case report

Shingo Kawano, Koichi Sato, Hiroshi Maekawa, Mutsumi Sakurada, Hajime Orita, Ryo Wada

ABSTRACT

Introduction: Primary duodenal adenocarcinoma is extremely rare. If this carcinoma occurs in fourth portion, it can now be diagnosed by recent developments in enteroscopy.

Case Report: We report a rare case of primary duodenal adenocarcinoma of the fourth portion diagnosed by double-balloon enteroscopy and resected surgically. A 57-year-old male was anemic. Positron emission tomography –computed tomography revealed accumulation in the fourth portion of the duodenum. Double-balloon enteroscope showed circular tumor of the fourth portion of the duodenum, and biopsy disclosed poorly differentiated adenocarcinoma. Partial duodenectomy and partial colonectomy were performed. The marginal artery of the transverse colon was invaded. Histological examination disclosed that the tumor was poorly differentiated adenocarcinoma and two lymph node metastases were seen.

Conclusion: Primary duodenal adenocarcinoma of fourth portion can be diagnosed by double-balloon enteroscopy and treated by surgical resection.

International Journal of Case Reports and Images (IJCRI)

International Journal of Case Reports and Images (IJCRI) is an international, peer reviewed, monthly, open access, online journal, publishing high-quality, articles in all areas of basic medical sciences and clinical specialties.

Aim of IJCRI is to encourage the publication of new information by providing a platform for reporting of unique, unusual and rare cases which enhance understanding of disease process, its diagnosis, management and clinico-pathologic correlations.

IJCRI publishes Review Articles, Case Series, Case Reports, Case in Images, Clinical Images and Letters to Editor.

Website: www.ijcasereportsandimages.com

CASE REPORT

OPEN ACCESS

Primary duodenal adenocarcinoma of the fourth portion diagnosed using double-balloon enteroscopy and surgically resected: A case report

Shingo Kawano, Koichi Sato, Hiroshi Maekawa, Mutsumi Sakurada, Hajime Orita, Ryo Wada

ABSTRACT

Introduction: Primary duodenal adenocarcinoma is extremely rare. If this carcinoma occurs in fourth portion, it can now be diagnosed by recent developments in enteroscopy. **Case Report:** We report a rare case of primary duodenal adenocarcinoma of the fourth portion diagnosed by double-balloon enteroscopy and resected surgically. A 57-year-old male was anemic. Positron emission tomography – computed tomography revealed accumulation in the fourth portion of the duodenum. Double-balloon enteroscope showed circular tumor of the fourth portion of the duodenum, and biopsy disclosed poorly differentiated adenocarcinoma. Partial duodenectomy and partial colectomy were performed. The marginal artery of the transverse colon was invaded. Histological examination disclosed that the tumor was poorly differentiated adenocarcinoma and two lymph node metastases were seen. **Conclusion:** Primary

duodenal adenocarcinoma of fourth portion can be diagnosed by double-balloon enteroscopy and treated by surgical resection.

Keywords: Double-balloon enteroscopy, Primary duodenal adenocarcinoma, The fourth portion

How to cite this article

Kawano S, Sato K, Maekawa M, Sakurada M, Orita H, Wada R. Primary duodenal adenocarcinoma of the fourth portion diagnosed using double-balloon enteroscopy and surgically resected: A case report. Int J Case Rep Images 2014;5(7):496–500.

doi:10.5348/ijcri-201487-CR-10398

INTRODUCTION

Primary duodenal adenocarcinoma is extremely rare, accounting for 0.3~0.4% of all gastrointestinal cancers [1]. It is too difficult to anatomically diagnose primary duodenal adenocarcinoma of the fourth portion. However, primary duodenal adenocarcinoma of the fourth portion can be diagnosed by recent developments in enteroscopy. Herein, we report a case of duodenal adenocarcinoma of the fourth portion diagnosed by double-balloon enteroscopy and resected surgically, and discuss it based on a review of literature.

CASE REPORT

A 57-year-old Japanese male was admitted to our hospital in December 2009 for evaluation of severe anemia. Past medical history included a gastric ulcer when he was 50-years-old. His family history was non-contributory. Physical examination on admission revealed

Shingo Kawano¹, Koichi Sato¹, Hiroshi Maekawa¹, Mutsumi Sakurada¹, Hajime Orita¹, Ryo Wada²

Affiliations: ¹Department of Surgery, Juntendo Shizuoka Hospital, Juntendo University School of Medicine, 1129 Nagaoka, Izunokuni-shi, Shizuoka 410-2295, Japan;

²Department of Pathology, Juntendo Shizuoka Hospital, Juntendo University School of Medicine, 1129 Nagaoka, Izunokuni-shi, Shizuoka 410-2295, Japan.

Corresponding Author: Shingo Kawano, Department of Surgery, Juntendo Shizuoka Hospital, Juntendo University School of Medicine, 1129 Nagaoka, Izunokuni-shi, Shizuoka 410-2295, Japan; Ph: +81-55-948-3111; Fax: +81-55-946-0514; Email: shingoshingo2000jp@yahoo.co.jp

Received: 25 October 2011

Accepted: 17 January 2012

Published: 01 July 2014

a body height 175 cm, weight 63 kg, blood pressure 112/61 mmHg, a regular pulse of 85/min, and body temperature 36.8°C. There was no sign of lymphadenopathy. His abdomen was soft and flat, and no abdominal, liver, or spleen masses were palpable.

There were no abnormal laboratory findings, except evidence of anemia (hemoglobin 6.6 g/dL) and an inflammatory reaction (white blood cell count was 8900/ μ g, and C-reactive protein was 3.7 mg/dL). Tumor markers, carcinoembryonic antigen (CEA) and carbohydrate antigen (CA19-9), were within normal ranges.

Abdominal computed tomography (CT) scan indicated a circumferential, thick, and unequally enhanced wall in the fourth portion of the duodenum, and showed no lymph node swelling or ascites. Upper gastrointestinal series demonstrated a 7-cm 'apple core' sign in the fourth portion of the duodenum (Figure 1). Double-balloon enteroscopy showed a circumferential tumor with ulceration in the fourth portion of the duodenum (Figure 2). The enteroscope could not pass the lesion. Biopsy specimen from the lesion disclosed poorly differentiated adenocarcinoma. Positron emission tomography (PET) scan revealed accumulation in the fourth portion of the duodenum (Figure 3). Based on these findings, the patient was diagnosed with primary duodenal adenocarcinoma of the fourth portion and underwent partial duodenectomy and partial colectomy because the marginal artery of the transverse colon was invaded by the tumor on January 22nd, 2010.

The resected tumor had deep ulceration with a round wall and measured 100×80 mm in size (Figure 4).

Histological examination disclosed that the tumor was poorly differentiated adenocarcinoma which invaded into the subserosal layer and two lymph node metastases were seen (Figure 5).

The patient had an uneventful postoperative course, and was discharged from hospital 40 days postoperatively. Combined chemotherapy of TS-1 and cisplatin was prescribed as neoadjuvant chemotherapy. TS-1 (80 mg/

Figure 2: Double-balloon enteroscopy: A circumferential tumor with ulceration in the fourth portion of the duodenum.

Figure 3: Positron Emission Tomography (PET): There was accumulation in the fourth portion of the duodenum.

Figure 1: Upper gastrointestinal series: A 7-cm 'apple core' sign was found in the fourth portion of the duodenum.

Figure 4: Surgical specimen: The resected tumor displayed deep ulceration with a round wall and measured 100×80 mm in size.

Figure 5: Histological findings: Poorly differentiated adenocarcinoma had invaded the subserosal layer. Two lymph node metastases were also detected.

m²) was administered 21 days, followed by 14 days rest as one course. Cisplatin (60 mg/m²) was administered on the eight day. Six courses of this adjuvant chemotherapy were administered. The patient is alive with no recurrence for one year.

DISCUSSION

Primary duodenal adenocarcinoma is extremely rare. Most primary duodenal adenocarcinomas are in the first and second portions, with 20% in the third portion, and 10% in the fourth portion [2]. Recently, reports of primary duodenal adenocarcinoma have been increasing because of progresses in gastric endoscopy [3, 4]. However, it is impossible to diagnose primary duodenal adenocarcinoma of the third and fourth portions by gastric endoscopy. There are still many cases to be diagnosed by the upper gastrointestinal series. In this case, primary duodenal adenocarcinoma of the fourth portion was diagnosed by double-balloon endoscopy. In Japan, there are no case reports of primary duodenal adenocarcinoma of the fourth portion diagnosed by double-balloon endoscopy. In Brazil, there is only one case report [5].

In Japan, there are eleven case reports of primary duodenal adenocarcinoma of the fourth portion resected surgically, including our case [5–16] (Table 1). The median age was 60.5 years (range 42–74 years). Six cases were men. Two cases invaded into the mucosal layer, three cases into the subserosal layer, and six cases went beyond the serosal layer. Seven cases had metastasis of the lymph nodes. According to these reports, there are two kinds of surgery, three cases of pylorus preserving pancreatoduodenectomy and partial

Table 1: Cases of the primary adenocarcinoma of the fourth portion of the duodenum reported in Japan

Author	Year	Age/Sex	Operation	Depth*	N	Prognosis
Adachi	1991	55/F	SD	SE	+	Unknown
Umehara	1996	42/F	PPPD	SE	+	24 mth. dead
Oshiro	2000	51/F	SD	SE	-	24 mth. alive
Saito	2001	54/M	SD	SE	+	Unknown
Hosokaewa	2003	74/F	SD	M	-	24 mth. alive
Maeda	2004	58/F	PPPD	SS	+	16 mth. alive
Suzuki	2005	66/M	PPPD	SE	+	24 mth. alive
Kinoshita	2006	65/M	SD	SS	-	11 mth. alive
Miyawaki	2006	72/M	SD	SE	+	30 mth. alive
Ishizaki	2008	72/M	SD	M	-	30 mth. alive
Our case	2010	57/M	SD	SS	+	12 mth. alive

PPPD - pylorus-preserving pancreatoduodenectomy, DS - segmental duodenectomy,

* Depth was expressed according to Japanese Classification of Gastric Carcinoma (The 14th edition), mth.: Month

duodenectomy and eight cases of partial duodenectomy with lymphadenectomy. In Kaklamanos's report, there was no significant difference in prognosis between pancreatoduodenectomy and partial duodenectomy in 63 cases of primary duodenal adenocarcinoma [16]. In Lowell's report, the seven cases of primary duodenal adenocarcinoma of the third or fourth portions are all alive after five years, except for those who died of another disease [17]. Five of seven cases underwent partial duodenectomy. There are many reports that partial duodenectomy is better in the case of primary duodenal adenocarcinoma in the third or fourth portions. Consequently, partial duodenectomy was thought to be a better surgery for primary distal duodenal adenocarcinoma. However, in Suzuki's report, a case of primary duodenal adenocarcinoma of the fourth portion underwent pancreatoduodenectomy [12]. There were metastases of the lymph nodes in No. 13 and No. 14 according to the Japanese classification of gastric carcinoma (The 14th Edition). That report insisted on the necessity of pancreatoduodenectomy with lymphadenectomy. Our case underwent partial duodenectomy with lymphadenectomy. There was no recurrence for one year in spite of advanced carcinoma and metastasis of the lymph nodes.

CONCLUSION

We reported an extremely rare case of primary duodenal adenocarcinoma of the fourth portion diagnosed by double-balloon enteroscopy and resected surgically.

Author Contributions

Shingo Kawano – Substantial contributions to conception and design, Acquisition of data, Analysis and interpretation of data, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Koichi Sato – Acquisition of data, Revising it critically for important intellectual content, Final approval of the version to be published

Hiroshi Maekawa – Acquisition of data, Revising it critically for important intellectual content, Final approval of the version to be published

Mutsumi Sakurada – Acquisition of data, Revising it critically for important intellectual content, Final approval of the version to be published

Hajime Orita – Acquisition of data, Revising it critically for important intellectual content, Final approval of the version to be published

Ryo Wada – Acquisition of data, Revising it critically for important intellectual content, Final approval of the version to be published

Guarantor

The corresponding author is the guarantor of submission.

Conflict of Interest

Authors declare no conflict of interest.

Copyright

© 2014 Shingo Kawano et al. This article is distributed under the terms of Creative Commons Attribution License which permits unrestricted use, distribution and reproduction in any medium provided the original author(s) and original publisher are properly credited. Please see the copyright policy on the journal website for more information.

REFERENCES

1. Tocci A, Mazzone G, Puma F, et al. Adenocarcinoma of the third and fourth portions of the duodenum: results of surgical treatment. *Arch Surg* 2003;138:80–5.
2. Shunsuke O, Gakuto K, Yoshihisa S, et al. A Clinicopathological Study of Our 10 Cases with Primary Carcinoma of the Duodenum. *The Japanese Journal of Gastroenterological Surgery* 2006;39:1458–63.
3. Rose DM, Hochwald SN, Klimstra DS, Brennan MF. Primary duodenal adenocarcinoma: a ten-year experience with 79 patients. *J Am Coll Surg* 1996;183:89–6.
4. Santoro E, Sacchi M, Scutari F, Carboni F, Graziano F. Primary adenocarcinoma of the duodenum: treatment and survival in 89 patients. *Hepatogastroenterology* 1997;44:1157–63.
5. Safatle-Ribeiro AV, Franzini TA, Kuqa R, et al. Double-balloon enteroscopy in the diagnosis of an adenocarcinoma of the fourth portion of the duodenum: report of a case. *Clinics (Sao Paulo)* 2007 Jun;62(3):353–6.
6. Keiichi A, Shiro F, Noriyuki A. A case of an advanced duodenal cancer of the fourth portion. *Clinics in Bowel Diseases* 1991;4:24–7.
7. Yasuhiko U, Tadatoshi O, Yoshihiko S, Toshio N, Naomi W, Ryutaro M. A case of an advanced duodenal cancer of the fourth portion. *Surgical Diagnosis & Treatment* 1996;38:1337–40.
8. Yoshio O, Kazuo S, Fuminori A, et al. A case of duodenal adenocarcinoma of the fourth portion diagnosed. *The Journal of the Hiroshima Medical Association* 2000;53:414–7.
9. Shuji S, Itaru E, Takafumi K, et al. A case of curatively resected duodenal cancer of the fourth portion. *The Japanese Journal of Gastroenterological Surgery* 2001;34:485–9.
10. Yukako H, Shigehiro K, Kimiyasu Y, et al. A case of duodenal polypoid intramucosal adenocarcinoma observed for six years. *Gastroenterological endoscopy* 2003;45:1130–4.
11. Tetsuya M, Akira T, Hironori K. A Case Report of Primary Duodenal Carcinoma of the Fourth Portion. *Journal of Japanese College of Surgeons* 2004;29:884–9.

12. Ryo S, Norichika M, Kentaro N, Akimasa Y, Takahisa M, Tomoaki M. A Case of Primary Adenocarcinoma of the Fourth Portion of the Duodenum Curatively Resected by Pylorus-Preserving Pancreatoduodenectomy. *The Japanese Journal of Gastroenterological Surgery* 2005;38:429–34.
13. Hiroyuki K, Katsutoshi T, Takeshi T, et al. A Resected Case of Primary Adenocarcinoma of the Fourth Portion of the Duodenum. *The Japanese Journal of Gastroenterological Surgery* 2006;39:1380–4.
14. Kiichiro M, Koichi T, Yu N, Daisuke K, Mitumasa A, Shinji T. A case of duodenal cancer of the fourth portion detected by routine gastroduodenoscopy. *Gastroenterological endoscopy* 2006;48:2639–44.
15. Tetsuo I, Makoto T, Takashi O, et al. A case of early cancer of the fourth portion of the duodenum. *The Japanese Journal of Gastroenterological Surgery* 2008;41:516–20.
16. Kaklamanos IG, Bathe OF, Franceschi D, Camarda C, Levi J, Livingstone AS. Extent of resection in the management of duodenal adenocarcinoma. *Am J Surg* 2000;179:37–41.
17. Lowell JA, Rossi RL, Munson JL, Braasch JW. Primary adenocarcinoma of third and fourth portions of duodenum, favorable prognosis after resection. *Arch Surg* 1992;127:557–60.

Access full text article on
other devices

Access PDF of article on
other devices

Edorium Journals: An introduction

Edorium Journals Team

About Edorium Journals

Edorium Journals is a publisher of high-quality, open access, international scholarly journals covering subjects in basic sciences and clinical specialties and subspecialties.

Invitation for article submission

We sincerely invite you to submit your valuable research for publication to Edorium Journals.

But why should you publish with Edorium Journals?

In less than 10 words - we give you what no one does.

Vision of being the best

We have the vision of making our journals the best and the most authoritative journals in their respective specialties. We are working towards this goal every day of every week of every month of every year.

Exceptional services

We care for you, your work and your time. Our efficient, personalized and courteous services are a testimony to this.

Editorial Review

All manuscripts submitted to Edorium Journals undergo pre-processing review, first editorial review, peer review, second editorial review and finally third editorial review.

Peer Review

All manuscripts submitted to Edorium Journals undergo anonymous, double-blind, external peer review.

Early View version

Early View version of your manuscript will be published in the journal within 72 hours of final acceptance.

Manuscript status

From submission to publication of your article you will get regular updates (minimum six times) about status of your manuscripts directly in your email.

Our Commitment

Six weeks

You will get first decision on your manuscript within six weeks (42 days) of submission. If we fail to honor this by even one day, we will publish your manuscript free of charge.

Four weeks

After we receive page proofs, your manuscript will be published in the journal within four weeks (31 days). If we fail to honor this by even one day, we will publish your manuscript free of charge and refund you the full article publication charges you paid for your manuscript.

Mentored Review Articles (MRA)

Our academic program "Mentored Review Article" (MRA) gives you a unique opportunity to publish papers under mentorship of international faculty. These articles are published free of charges.

Most Favored Author program

Join this program and publish any number of articles free of charge for one to five years.

Favored Author program

One email is all it takes to become our favored author. You will not only get fee waivers but also get information and insights about scholarly publishing.

Institutional Membership program

Join our Institutional Memberships program and help scholars from your institute make their research accessible to all and save thousands of dollars in fees make their research accessible to all.

Our presence

We have some of the best designed publication formats. Our websites are very user friendly and enable you to do your work very easily with no hassle.

Something more...

We request you to have a look at our website to know more about us and our services.

We welcome you to interact with us, share with us, join us and of course publish with us.

Edorium Journals: On Web

Browse Journals

CONNECT WITH US