Early bilateral hippocampal lesions in transient global amnesia: Evidence against delayed ischemia?

Kelsey Flynn, Pascale Lavoie, Robert Laforce Jr

ABSTRACT

Abstract is not required for Clinical Images

All Articles Open Access

International Journal of Case Reports and Images (IJCRI)

International Journal of Case Reports and Images (IJCRI) is an international, peer reviewed, monthly, open access, online journal, publishing high-quality, articles in all areas of basic medical sciences and clinical specialties.

Aim of IJCRI is to encourage the publication of new information by providing a platform for reporting of unique, unusual and rare cases which enhance understanding of disease process, its diagnosis, management and clinico-pathologic correlations.

IJCRI publishes Review Articles, Case Series, Case Reports, Case in Images, Clinical Images and Letters to Editor.

Website: www.ijcasereportsandimages.com

CLINICAL IMAGES

OPEN ACCESS

Early bilateral hippocampal lesions in transient global amnesia: Evidence against delayed ischemia?

Kelsey Flynn, Pascale Lavoie, Robert Laforce Jr

CASE REPORT

A 65-year-old right-handed female was admitted to neurology after she suddenly began experiencing problems with her memory. Her past medical history was significant for osteoarthritis and a fractured clavicle, but there were no vascular risk factors, head trauma, migraine or epilepsy. She did not smoke, occasionally drank alcohol and denied illicit drug use. Her mother had passed away at 98 years of age without any sign of Alzheimer's disease but one of her aunts was diagnosed with Alzheimer's disease. She has four siblings, all were in good health, except for one brother with cardiac problems.

On the day of the event, at 11:15 a.m. she began showing symptoms after a short bout of intense running for almost six minutes. She reported a vague sensation of confusion as to where she was and what she was doing, but nonetheless called her husband to pick her up. When he arrived, he noted her confusion in space and time, and she admitted to feeling dizzy. Upon arrival at their home, her confusion persisted. For example, she had forgotten about her sister's birthday, and that they were renovating their kitchen. She repeatedly asked her husband "Why is there a present on the counter?", or "Why is our kitchen such a mess?" After a few hours, she returned to normal level of functioning with no recollection of what had just happened.

Neurological examination was normal. Blood pressure was 122/68 mmHg, heart rate 76 bpm. Basic blood work, ions, urea/creatinine, TSH, calcium/magnesium/phosphorus, B12/folates were unremarkable. She scored 26/30 on a brief cognitive screening measure, losing four points in the free recall section, but showing intact recognition with verbal cues. The rest of her cognitive examination was normal. Magnetic resonance imaging (MRI) scan of the brain conducted 19 hours after symptom onset revealed bilateral lesions of the hippocampi (Figure 1).

DISCUSSION

The patient was diagnosed with transient global amnesia (TGA), a benign syndrome characterized by the sudden onset of severe anterograde and mild retrograde amnesia [1]. Transient global amnesia is generally accompanied by repetitive questioning and resolves within 24 hours, leaving no sequelae. Recurrence is rare. Several possible etiologies have been suggested as underlying mechanisms for TGA including migraine-related mechanisms, venous-flow abnormalities, epileptic seizures, and ischemic events [2–4]. However,

Figure 1: Magnetic resonance imaging scan of the brain (A) Showing bilateral hyperintense posterior portions of the hippocampi, (B) Hypointense apparent diffusion coefficient map in similar brain loci, both compatible with ischemic lesions. Follow-up Magnetic resonance imaging conducted six weeks later showed complete resolution of the lesions.

Kelsey Flynn¹, Pascale Lavoie², Robert Laforce Jr³

Affiliations: ¹BSc, University of British Columbia, Vancouver, Canada; ²MD, Faculté de Médecine, Université Laval, Québec, Canada; ³MD, PhD, Faculté de Médecine, Université Laval, Québec, Canada; Clinique Interdisciplinaire de Mémoire, Département des Sciences Neurologiques, Hôpital de l'Enfant-Jésus, Québec, Canada.

Corresponding Author: Robert Laforce Jr, MD, PhD, Clinique Interdisciplinaire de Mémoire, CHU de Québec, 1401, 18ième rue, Québec, Canada, G1J 1Z4; Tel: 418-649-5980, Fax: 418-649-5981; Email: robert.laforce@fmed.ulaval.ca

Received: 31 July 2014
Accepted: 14 August 2014
Published: 01 November 2014

the fundamental etiology of TGA remains unclear and may be highly heterogeneous.

Recent neuroimaging studies have revealed that a large portion of individuals with TGA present with small, reversible, restricted lesions on diffusion MRI [5]. This has been used to support an underlying ischemic event that could be the result of venous overflow. Furthermore, these lesions often tend to appear when patients are scanned after a 48-hour delay, hence supporting the delayed ischemic hypothesis [5]. Interestingly, our patient's ischemic event was detected in the acute phase (0–24 hours) of TGA where most other authors have failed to report any changes. Whether this is attributable to lesion size is unclear, but nonetheless does not support the notion of delayed ischemia.

We herein report a case of transient global amnesia (TGA) secondary to bilateral ischemic hippocampal lesions 19 hours after symptom onset. In accord with the hypothesized etiology of venous overflow, our findings replicate other studies showing reversible ischemia in hippocampal regions associated with this syndrome. Interestingly, lesions to other structures known to be involved in declarative memory (e.g., thalamus, cingulate gyrus, and basal ganglia) have been associated with TGA [6]. In addition to being rare because of their bilateral nature, the lesions shown here are unique in that very few authors have reported such changes in the acute phase of TGA. Moreover, they raise the possibility that delayed ischemia may not be the sole mechanism behind TGA. Indeed, ischemia may be present early in the process but simply not detected [5].

CONCLUSION

Newly developed imaging techniques such as functional Magnetic resonance imaging scan may allow earlier detection of cerebral changes associated with transient global amnesia. This in turn may help clinicians in the differential diagnosis of other transient amnesic syndromes, as well as more rapid targeted interventions.

How to cite this article

Flynn K, Lavoie P, Laforce R Jr. Early bilateral hippocampal lesions in transient global amnesia: Evidence against delayed ischemia? Int J Case Rep Images 2014;5(11):795–797.

doi:10.5348/ijcri-201467-CL-10055

Acknowledgements

We would like to acknowledge the patient who kindly allowed us to publish her history.

Author Contributions

Kelsey Flynn – Analysis and interpretation of data, Drafting the article, Final approval of the version to be published

Pascale Lavoie – Acquisition of data, Analysis and interpretation of data, Critical revision of the article, Final approval of the version to be published

Robert Laforce Jr – Conception and design, Acquisition of data, Analysis and interpretation of data, Critical revision of the article, Final approval of the version to be published

Guarantor

The corresponding author is the guarantor of submission.

Conflict of Interest

Authors declare no conflict of interest.

Copyright

© 2014 Kelsey Flynn et al. This article is distributed under the terms of Creative Commons Attribution License which permits unrestricted use, distribution and reproduction in any medium provided the original author(s) and original publisher are properly credited. Please see the copyright policy on the journal website for more information.

REFERENCES

1. Bartsch T, Deuschl G. Transient Global Amnesia: Functional anatomy and clinical implications. *Lancet Neurol* 2010 Feb;9(2):205–14.
2. Quinette P, Guillery-Girard B, Dayan J, et al. What does transient global amnesia really mean? Review of the literature and thorough study of 142 cases. *Brain* 2006 Jul;129(Pt 7):1640–58.
3. Olsen J, Jørgensen MB. Leao's spreading depression in the hippocampus explains transient global amnesia. A hypothesis. *Acta Neurol Scand* 1986 Feb;73(2):219–20.
4. Lewis SL. Aetiology of Transient Global Amnesia. *Lancet* 1998 Aug 1;352(9125):397–9.
5. Sedlaczek O, Hirsch JG, Grips E, et al. Detection of Delayed Focal MR Changes in the Lateral Hippocampus in Transient Global Amnesia. *Neurology* 2004 Jun 22;62(12):2165–70.
6. Gallardo-Tur A, Romero-Godoy J, de la Cruz Cosme C, Arboix A. Transient Global Amnesia Associated with an Acute Infarction at the Cingulate Gyrus. *Case Rep Neurol Med* 2014;2014:418180.

Access full text article on
other devices

Access PDF of article on
other devices

Edorium Journals: An introduction

Edorium Journals Team

About Edorium Journals

Edorium Journals is a publisher of high-quality, open access, international scholarly journals covering subjects in basic sciences and clinical specialties and subspecialties.

Invitation for article submission

We sincerely invite you to submit your valuable research for publication to Edorium Journals.

But why should you publish with Edorium Journals?

In less than 10 words - we give you what no one does.

Vision of being the best

We have the vision of making our journals the best and the most authoritative journals in their respective specialties. We are working towards this goal every day of every week of every month of every year.

Exceptional services

We care for you, your work and your time. Our efficient, personalized and courteous services are a testimony to this.

Editorial Review

All manuscripts submitted to Edorium Journals undergo pre-processing review, first editorial review, peer review, second editorial review and finally third editorial review.

Peer Review

All manuscripts submitted to Edorium Journals undergo anonymous, double-blind, external peer review.

Early View version

Early View version of your manuscript will be published in the journal within 72 hours of final acceptance.

Manuscript status

From submission to publication of your article you will get regular updates (minimum six times) about status of your manuscripts directly in your email.

Our Commitment

Six weeks

You will get first decision on your manuscript within six weeks (42 days) of submission. If we fail to honor this by even one day, we will publish your manuscript free of charge.

Four weeks

After we receive page proofs, your manuscript will be published in the journal within four weeks (31 days). If we fail to honor this by even one day, we will publish your manuscript free of charge and refund you the full article publication charges you paid for your manuscript.

Mentored Review Articles (MRA)

Our academic program "Mentored Review Article" (MRA) gives you a unique opportunity to publish papers under mentorship of international faculty. These articles are published free of charges.

Favored Author program

One email is all it takes to become our favored author. You will not only get fee waivers but also get information and insights about scholarly publishing.

Institutional Membership program

Join our Institutional Memberships program and help scholars from your institute make their research accessible to all and save thousands of dollars in fees make their research accessible to all.

Our presence

We have some of the best designed publication formats. Our websites are very user friendly and enable you to do your work very easily with no hassle.

Something more...

We request you to have a look at our website to know more about us and our services.

We welcome you to interact with us, share with us, join us and of course publish with us.

Edorium Journals: On Web

Browse Journals

CONNECT WITH US