Patent omphalomesenteric duct opening into the vermiform appendix

**Akintayo David OlaOlorun, Stephen Adesope Adesina,
Adepeju Olatayo Adepoju, Isaac Olusayo Amole**

ABSTRACT

Introduction: Persistent omphalomesenteric duct anomalies are very rare the world over. These anomalies are in a spectrum and present in various forms, ranging from the infamous Meckel's diverticulum to a fibrous non-patent connection between the umbilicus and the distal ileum, to a patent connection between the umbilicus and the intestine, the patent omphalomesenteric duct. Case Report: A two-month-old female child who was born with a membrane-covered umbilical mass which after sometime, became two swellings with intestinal epithelial covering and fistulous openings. At surgical exploration, she was discovered to have a patent omphalomesenteric duct which opened into the appendix, as well as intestinal malrotation abnormality. Conclusion: The occurrence of patent omphalomesenteric duct opening into the vermiform appendix is a rare occurrence and there should be high index of suspicion when there is failure of healing of the umbilical stump and there is discharge of intestinal contents via the umbilicus.

International Journal of Case Reports and Images (IJCRI)

International Journal of Case Reports and Images (IJCRI) is an international, peer reviewed, monthly, open access, online journal, publishing high-quality, articles in all areas of basic medical sciences and clinical specialties.

Aim of IJCRI is to encourage the publication of new information by providing a platform for reporting of unique, unusual and rare cases which enhance understanding of disease process, its diagnosis, management and clinico-pathologic correlations.

IJCRI publishes Review Articles, Case Series, Case Reports, Case in Images, Clinical Images and Letters to Editor.

Website: www.ijcasereportsandimages.com

CASE REPORT

PEER REVIEWED | OPEN ACCESS

Patent omphalomesenteric duct opening into the vermiform appendix

Akintayo David OlaOlorun, Stephen Adesope Adesina,
Adepeju Olatayo Adepoju, Isaac Olusayo Amole

ABSTRACT

Introduction: Persistent omphalomesenteric duct anomalies are very rare the world over. These anomalies are in a spectrum and present in various forms, ranging from the infamous Meckel's diverticulum to a fibrous non-patent connection between the umbilicus and the distal ileum, to a patent connection between the umbilicus and the intestine, the patent omphalomesenteric duct. **Case Report:** A two-month-old female child who was born with a membrane-covered umbilical mass which after sometime, became two swellings with intestinal epithelial covering and fistulous openings. At surgical exploration, she was discovered to have a patent omphalomesenteric duct which opened into the appendix, as well as intestinal malrotation abnormality. **Conclusion:** The occurrence of patent omphalomesenteric duct opening into the vermiform appendix is a rare occurrence and there should be high index of suspicion when there is failure of healing of the umbilical stump and there is discharge of intestinal contents via the umbilicus.

Akintayo David OlaOlorun¹, Stephen Adesope Adesina², Adepeju Olatayo Adepoju², Isaac Olusayo Amole³

Affiliations: ¹MBBS, FMCGP, Consultant, Family Medicine, Bowen University Teaching Hospital, Ogbomoso, Nigeria; ²MBBS, MWACP(FM), Senior Registrar, Family Medicine, Bowen University Teaching Hospital, Ogbomoso, Nigeria; ³MBBS, FWACP(FM), Consultant, Family Medicine, Bowen University Teaching Hospital, Ogbomoso, Nigeria.

Corresponding Author: Isaac Olusayo Amole, Bowen University Teaching Hospital, P. O. Box 15, Ogbomoso, Oyo State, Nigeria, 009234; Ph: +2348057078851; Email: amoleio@yahoo.com

Received: 04 September 2015

Accepted: 17 October 2015

Published: 01 December 2015

Keywords: Appendix, Carcinoid tumor, Embryonic vitelline duct, Intestinal malrotation

How to cite this article

OlaOlorun AD, Adesina SA, Adepoju AO, Amole IO. Patent omphalomesenteric duct opening into the vermiform appendix. Int J Case Rep Images 2015;6(12):743–746.

doi:10.5348/ijcri-2015119-CR-10580

INTRODUCTION

Persistent omphalomesenteric duct anomalies are very rare the world over. In a report out of Zaria, Nigeria, Ameh et al. saw less than one patient per year. They treated 18 patients in 22 years [1]. These anomalies are in a spectrum and present in various forms, ranging from the infamous Meckel's diverticulum to a fibrous non-patent connection between the umbilicus and the distal ileum, to a patent connection between the umbilicus and the intestine, the patent omphalomesenteric duct. It is normally observed as a connection with the distal ileum but has been known to connect with the cecum, ascending colon and the appendix [2, 3]. We present the case of a two-month-old female child who was referred to us with complaints of failure of healing of the umbilical stump and the discharge of intestinal contents via the umbilicus.

CASE REPORT

A two-month-old female child was referred to us with failure to heal her umbilical stump. She was born the second of a sibship of two, to parents in their thirties after an uneventful pregnancy and delivery. After delivery, her mother noticed that the child had an umbilical mass,

covered by a membrane. The umbilicus was ligated normally and the stump was held in a plastic umbilical clamp. Since this child was born at a prayer house, upon leaving there, her mom went to an orthodox hospital where the caregivers commenced daily dressing changes to the umbilicus. This continued for a while but when the parents and the grandmother noticed a lack of healing of the umbilicus at the time they expected it to have healed, they became concerned. Shortly afterwards, the mother noticed an intermittent discharge of what appeared to be intestinal contents from the umbilicus and this prompted the referral to our hospital.

Upon arrival, the child was examined closely for congenital abnormalities of the heart, spine and anus and there were none. The rest of her physical examinations were normal except for the umbilicus which had what looked like intestinal mucosa in two bumps (Figure 1A–B). There were fistulous openings on either side of the bumps but we could not express out yellowish-brownish fluid suggestive of intestinal contents. The clinical impression after assessing the child was that of a patent omphalomesenteric duct. Further radiological evaluation of this apparent umbilical fistula was not carried out because of the financial difficulties of the young family. The child underwent an umbilical stump exploration via a circum-umbilical incision.

Operative findings: The two bumps were found to be the tip of the appendix which was discovered to be attached to the umbilicus. There was associated malrotation of the intestine, with the cecum freely mobile, not attached to the posterior abdominal wall as expected, and it was located to the left side of the terminal ileum (Figure 2) as opposed to its normal anatomy. The blood supply to the appendix was found to be located anterior to the appendix, cecum and ileum in an aberrant position. The appendix was removed using the standard appendectomy technique, the fascia was closed and the umbilicus reconstructed (Figures 3 and 4). The child made an uneventful recovery. The post-operative diagnosis was patent omphalomesenteric duct which opened into the appendix.

DISCUSSION

The rarity of persistent omphalomesenteric duct remnants of all kinds [1, 4] is such that many physicians will go through their whole medical career without seeing one. It is accepted that abnormal umbilical masses and/or fistulas represent a failure of obliteration and delay of involution of the embryonic vitelline (omphalomesenteric) duct. The most frequently seen of the aforementioned spectrum of findings is the Meckel's diverticulum and it is reported to be present in only 2–3% of people [5], and to be more symptomatic in males. The appendico-umbilical fistula is believed to be a variant of the patent omphalomesenteric duct, one that connected with the appendix instead of the distal ileum or other

places like the cecum or the ascending colon [2], and that indeed is the diagnosis in this case.

When an umbilical appendix is present, it is known to be associated with various forms of malrotation of the gut [3, 6]. The aberrant position of the appendiceal blood vessels in this case supports the diagnosis of associated malrotation. The significance of this is that these children are susceptible to the well-known potential complications of malrotation such as midgut volvulus. There can also be twisting of the gut around the attachment of the appendix to the umbilicus. It is imperative to remove the umbilical

Figure 1: (A) Umbilical “bumps” clearly showing intestinal mucosa, (B) Umbilical “bumps”, lateral view.

Figure 2: Malrotation of the gut with the cecum freely mobile and to the left of the terminal ileum

Figure 3: Immediate postoperative umbilical reconstruction

Figure 4: Postoperative (four weeks) picture of the umbilicus.

appendix (or the persisting omphalomesenteric duct) as soon as possible to prevent this complication because of this. Other complications that can occur in persons with Meckel's diverticulum are hemorrhage and growth of Carcinoid tumor [5] and presumably this can also occur when the patent omphalomesenteric duct opens into the appendix instead of the distal ileum. Obviously, having had her omphalomesenteric duct/appendix removed, the child in this case has had these potential complications forestalled. The preoperative diagnosis of cases like this can be clinically difficult and once operated, the procedure chosen to deal with the problem has to be individualized.

CONCLUSION

The occurrence of patent omphalomesenteric duct opening into the vermiform appendix is a rare occurrence and there should be high index of suspicion when there is failure of healing of the umbilical stump and there is discharge of intestinal contents via the umbilicus.

Author Contributions

Akintayo David OlaOlorun – Substantial contributions to conception and design, Drafting the article, Revising it critically for important intellectual content, Final approval of the version to be published

Stephen Adesope Adesina – Acquisition of data, Revising it critically for important intellectual content, Final approval of the version to be published

Adepeju O. Adegoke – Acquisition of data, Revising it critically for important intellectual content, Final approval of the version to be published

Isaac Olusayo Amole – Acquisition of data, Critical revision of the article, Final approval of the version to be published

Guarantor

The corresponding author is the guarantor of submission.

Conflict of Interest

Authors declare no conflict of interest.

Copyright

© 2015 Akintayo David OlaOlorun et al. This article is distributed under the terms of Creative Commons Attribution License which permits unrestricted use, distribution and reproduction in any medium provided the original author(s) and original publisher are properly credited. Please see the copyright policy on the journal website for more information.

REFERENCES

1. Ameh EA, Mshelbwala PM, Dauda MM, Sabiu L, Nmadu PT. Symptomatic vitelline duct anomalies in children. *S Afr J Surg* 2005 Aug;43(3):84–5.
2. Crankson SJ, Ahmed GS, Palkar V. Patent omphalomesenteric duct of the vermiform appendix in a neonate: congenital appendicoumbilical fistula. *Pediatr Surg Int* 1998 Dec;14(3):229–30.
3. Fuijkschoot J, Wijnen RM, Gerrits GP, Dubois SV, Rieu PN. A neonate with an intact congenital umbilical appendix: an alternative theory on the etiology of the appendico-umbilical fistula. *Pediatr Surg Int* 2006 Aug;22(8):689–93.
4. Collins DC. A study of 50,000 specimens of the human vermiform appendix. *Surg Gynecol Obstet* 1955 Oct;101(4):437–45.
5. Dumper J, Mackenzie S, Mitchell P, Sutherland F, Quan ML, Mew D. Complications of Meckel's diverticula in adults. *Can J Surg* 2006 Oct;49(5):353–7.
6. Cevik M, Boleken ME, Kadioglu E. Appendicoumbilical fistula: a rare reason for neonatal umbilical mass. *Case Rep Med* 2011;2011:835474.

Access full text article on
other devices

Access PDF of article on
other devices

Edorium Journals: An introduction

Edorium Journals Team

About Edorium Journals

Edorium Journals is a publisher of high-quality, open access, international scholarly journals covering subjects in basic sciences and clinical specialties and subspecialties.

Invitation for article submission

We sincerely invite you to submit your valuable research for publication to Edorium Journals.

But why should you publish with Edorium Journals?

In less than 10 words - we give you what no one does.

Vision of being the best

We have the vision of making our journals the best and the most authoritative journals in their respective specialties. We are working towards this goal every day of every week of every month of every year.

Exceptional services

We care for you, your work and your time. Our efficient, personalized and courteous services are a testimony to this.

Editorial Review

All manuscripts submitted to Edorium Journals undergo pre-processing review, first editorial review, peer review, second editorial review and finally third editorial review.

Peer Review

All manuscripts submitted to Edorium Journals undergo anonymous, double-blind, external peer review.

Early View version

Early View version of your manuscript will be published in the journal within 72 hours of final acceptance.

Manuscript status

From submission to publication of your article you will get regular updates (minimum six times) about status of your manuscripts directly in your email.

Our Commitment

Six weeks

You will get first decision on your manuscript within six weeks (42 days) of submission. If we fail to honor this by even one day, we will publish your manuscript free of charge.

Four weeks

After we receive page proofs, your manuscript will be published in the journal within four weeks (31 days). If we fail to honor this by even one day, we will publish your manuscript free of charge and refund you the full article publication charges you paid for your manuscript.

Most Favored Author program

Join this program and publish any number of articles free of charge for one to five years.

Favored Author program

One email is all it takes to become our favored author. You will not only get fee waivers but also get information and insights about scholarly publishing.

Institutional Membership program

Join our Institutional Memberships program and help scholars from your institute make their research accessible to all and save thousands of dollars in fees make their research accessible to all.

Our presence

We have some of the best designed publication formats. Our websites are very user friendly and enable you to do your work very easily with no hassle.

Something more...

We request you to have a look at our website to know more about us and our services.

We welcome you to interact with us, share with us, join us and of course publish with us.

Edorium Journals: On Web

Browse Journals

CONNECT WITH US